



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2001

Nutrition and prostate cancer

Schmitz-Dräger, B J ; Eichholzer, Monika ; Beiche, B ; Ebert, T

Abstract: Nutrition is apparently a major risk factor for the development and progression of prostate cancer. Based on experimental studies and epidemiologic data mainly from case-control studies or cohort studies, there is strong evidence that reduction of the total energy consumption, a diet comprising less than 30% fat, and increased intake of phytoestrogens, vitamins D and E and selenium could yield a decreased prostate cancer incidence. Furthermore, some of these measures appear to have antitumoral capacity even in the presence of the disease. These observations have provided a rationale to forward large prospective trials on dietary interventions to prove the efficacy of the concept and further delineate the correlation between nutritional compounds and prostate cancer risk. These chemoprevention trials are either aiming a reduction prostate cancer incidence or a decrease in tumor progression. Depending on the study design, large numbers of individuals need to be enrolled and long follow-up intervals are required thus making such trials highly complex and cost-intensive. However, regarding the potential relevance of chemoprevention on public health, further efforts to identify nutritional factors affecting prostate cancer growth are warranted.

DOI: <https://doi.org/10.1159/000050935>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-109539>

Journal Article

Published Version

Originally published at:

Schmitz-Dräger, B J; Eichholzer, Monika; Beiche, B; Ebert, T (2001). Nutrition and prostate cancer. *Urologia internationalis*, 67(1):1-11.

DOI: <https://doi.org/10.1159/000050935>

Bernd J. Schmitz-Dräger^a
Monika Eichholzer^b
Birgit Beiche^a
Thomas Ebert^a

^a Urology, EuromedClinic, Fürth, Germany,
and

^b Institute for Social and Preventive
Medicine, University of Zürich,
Switzerland

Nutrition and Prostate Cancer

Abstract

Nutrition is apparently a major risk factor for the development and progression of prostate cancer. Based on experimental studies and epidemiologic data mainly from case-control studies or cohort studies, there is strong evidence that reduction of the total energy consumption, a diet comprising less than 30% fat, and increased intake of phytoestrogens, vitamins D and E and selenium could yield a decreased prostate cancer incidence. Furthermore, some of these measures appear to have antitumoral capacity even in the presence of the disease. These observations have provided a rationale to forward large prospective trials on dietary interventions to prove the efficacy of the concept and further delineate the correlation between nutritional compounds and prostate cancer risk. These chemoprevention trials are either aiming a reduction prostate cancer incidence or a decrease in tumor progression. Depending on the study design, large numbers of individuals need to be enrolled and long follow-up intervals are required thus making such trials highly complex and cost-intensive. However, regarding the potential relevance of chemoprevention on public health, further efforts to identify nutritional factors affecting prostate cancer growth are warranted.

Introduction

Epidemiologic studies have demonstrated that the incidence of clinically overt prostate cancer is about 10 times higher in White US American men as compared to Japanese men with a comparable socioeconomic background [1, 2]. Furthermore, migration studies showed that the low prostate cancer incidence in Japanese immigrants in the US rose about 3 times within one generation. Also in Europe there is a significant difference in prostate cancer incidence from 11.8/100,000 male population in Portugal to 50.2/100,000 males in Sweden.

Although prostate cancer is one of the most common cancers in western Countries, its etiology is mostly unknown. However, aside of age and ethnicity, several environmental factors, including diet, have been identified in epidemiological studies, although these findings have not been completely consistent [3–5]. So far, several nutritional compounds have been suggested to influence prostate cancer (table 1). Through the past decade the understanding of the molecular mechanisms potentially underlying these effects have been identified thus now providing the basis for interventional studies.

KARGER

Fax + 41 61 306 12 34
E-Mail karger@karger.ch
www.karger.com

© 2001 S. Karger AG, Basel
0042–1138/01/0671–00010/\$17.50

Accessible online at:
www.karger.com/journals/uin

B.J. Schmitz-Dräger
Urology, EuromedClinic
Europa-Allee 1, D–90763 Fürth (Germany)
Tel. +49 911 9714 531, Fax +49 911 9714 532
E-Mail bsd@euromed.de

Table 1. Nutritional compounds potentially involved in the development of prostate cancer

Phytoestrogens
Antioxidants (carotenoids, vitamins A, C, D, E)
Selenium
Fat/total energy uptake

Phytoestrogens

Testosterone is known to play a crucial role in the development and progression of benign and malignant prostatic disease. Although it remains a matter of discussion if testosterone serum levels are correlated with the incidence of prostate cancer [6–8], several observations suggest that free testosterone might be one of the factors inducing and stimulating prostate cancer. Phytoestrogens are known to be strong inhibitors of 5- α -reductase thus potentially counteracting testosterone-mediated stimulation of prostatic growth [9, 10].

There is some epidemiologic evidence that phytoestrogens may influence progression of benign and malignant prostatic disease [5, 11, 12]. Diets in areas with a low prostate cancer incidence (far-east Asia, southern Europe) are rich in fibers and vegetables containing high amounts of isoflavonoids, flavonoids and lignans, the most important categories of phytoestrogens. While high concentration of flavonoids and isoflavonoids, e.g. daidzein or genistein, are detected in vegetables (e.g. soy) and fruits, lignans can also be found at high concentrations in linseeds, grains and cereals. Phytoestrogens may influence growth and proliferation of prostate cells through different mechanisms (table 2). As expected, serum levels or urinary concentrations of phytoestrogen metabolites are much higher in men from areas with low prostate cancer incidence as compared to men from western countries [11–13].

Despite experimental and epidemiologic evidence from ecological studies on the beneficial effects of phytoestrogens, only few data from case-control and cohort studies have been published so far. Two case-control studies report a significant inverse correlation between phytoestrogen intake and prostate cancer risk [14, 15]. These observations are further supported by the summarized data from three case-control studies including 617 prostate cancer cases from Canada [16]. The intake of green vegetables, beans, lentils, nuts and cruciferous vegetables was inversely correlated with the incidence of prostate cancer.

Table 2. Putative mechanisms of phytoestrogen action on prostate cells

Increase of SHBG serum concentration and subsequent decrease of free testosterone through binding to liver estrogen receptors
Decrease of DNA synthesis through inhibition of tyrosine kinase and topoisomerase
Decrease of the effects of free radicals through antioxidant properties
Inhibition of cytochrome P450 activation
Neoangiogenesis inhibition
Inhibition of intraprostatic testosterone metabolism through inhibition of 5 α -reductase and aromatase

Similar results were obtained in two cohort studies from the US, both showing significant beneficial effects of soybean product consumption on prostate cancer risk [17, 18]. However, despite these encouraging data, the assessment of dietary phytoestrogen consumption is highly complex and the results of these trials may be confounded by multiple parameters [19]. Therefore, further assessment of this topic is required.

Lycopene

Lycopene is a carotenoid that is present in tomatoes, processed tomato products and several fruits. It is one of the most potent antioxidants among dietary carotenoids. Dietary intake of tomatoes and tomato products has been shown to be associated with a decreased risk of chronic diseases [20]. Although the antioxidant properties of lycopene are thought to be primarily responsible for its effects, evidence is accumulating that other mechanisms may also be involved.

The effects of lycopene on prostate cancer have been extensively addressed in a variety of studies. While case-control studies by Le Marchand et al. [21], Key et al. [22] and Hayes et al. [23] showed no differences between those individuals on a lycopene-rich diet and the controls, the trials by Norrish et al. [24] and Tzonou et al. [25] yielded a decreased risk of developing prostate cancer for individuals consuming high amounts of tomatoes or having a lycopene-rich diet reaching significance only in the latter trial (table 3).

Furthermore, the impact of lycopene on prostate cancer risk has also been studied in several cohort studies [26–31] (table 3). All trials studying the effects of tomato-rich diets showed a significant decrease of the

Table 3. Summary of trials investigating the influence of lycopene and tomatoes/tomato products on the incidence of prostate cancer

Reference	Cases	Measure	Ass.	RR/OR (95% CI) ¹
<i>Case-control trials</i>				
Le Marchand (1991) [21]	452	Diet	NS	0.9 <70 years
			NS	1.1 ≥70 years
Key (1997) [22]	328	Diet	NS	0.99 (0.68–1.45)
Tzonou (1997) [25]	320	Tomatoes	↓	0.70
Hayes (1999) [23]	932	Diet	NS	
Norrish (2000) [24]	317	Diet	NS	0.76 (0.50–1.17)
<i>Cohort studies or nested case-control studies</i>				
Mills (1989) [26]	180	Tomatoes	↓	0.60 (0.37–0.97)
Hsing (1990) [27]	103	Serum	NS	0.50 (0.20–1.29)
Giovannucci (1995) [28]	773	Tomato products	↓	0.65 (0.44–0.95)
Nomura (1997) [29]	142	Serum	NS	1.1 (0.5–2.2)
Cerhan (1997) [30]	1,050	Tomatoes	↓	0.5 (0.3–0.9)
Gann (1998) [31]	578	Plasma	↓	0.56 (0.34–0.92)

NS = Statistically not significant; ↓ = statistically significant inverse association.

¹ Comparing extreme categories, low = reference.

prostate cancer risk in individuals with a high consumption of tomatoes or tomato products [26, 28, 30]. However, it remains unclear if these beneficial effects are related to lycopene intake or rather to the multiple other nutritional compounds in tomatoes.

In three other trials, lycopene serum or plasma levels were studied [27, 29, 31]. While Nomura et al. [29] did not find a difference between the study groups, Gann et al. [31] showed a correlation with a significant decrease of the prostate cancer risk in the group with high lycopene plasma levels. A similar decrease was also seen by Hsing et al. [27], however, because of the small number of prostate cancer cases (n = 103), this observation was not significant.

In a recent review on 72 studies concerning the impact of tomatoes and tomato-based products on the incidence of various cancers, Giovannucci [32] identified 35 trials reporting a significant inverse relationship. The evidence for a benefit was strongest for cancers of the prostate, lung and stomach. Moreover, no study indicated that higher tomato consumption or lycopene blood levels statistically significantly increased the risk of cancer of any site.

Vitamins

From epidemiologic evaluations there is a great body of evidence of an inverse correlation between the intake of vitamins and the incidence of different types

of cancer. These observations are further supported by results of experimental trials. Specifically with regard to the development of prostate cancer, vitamins A, C, D and E have been suggested to be of relevance.

The molecular pathways through which vitamins may act are severalfold. Neutralization of the oxidative capacity of free radicals in the tissue appears to be a major way of action for several vitamins [33]. Furthermore, stimulation of the immune system and its putative antitumoral actions has been shown to be mediated through vitamins. Retinoids and vitamin A are known to be strong inducers of cellular differentiation. Within this context the loss of intercellular interaction, another aspect of dedifferentiation, can be restored under the influence of vitamins. Apart from these indirect actions, direct antiproliferative effects of several vitamins in vitro and in vivo have been reported [33–38].

Total Vitamin A (β-Carotene and Retinol)

There is some experimental evidence that the provitamins or vitamin A may influence the development and progression of prostate cancer. Williams et al. [38] demonstrated an in vitro down-regulation of the proliferation of prostate cancer cells in the presence of β-carotene. Further investigation suggests a metabolic conversion of β-carotene to retinol by the prostate tumor cells. This observation supports the results of a

Table 4. Summary of trials investigating the influence of vitamin C on the incidence or mortality of prostate cancer

Reference	Cases	Measure	Ass.	RR/OR (95% CI) ¹
<i>Case-control</i>				
Vlajinac (1997) [56]	101	Diet	NS	0.90 (0.48–1.69)
Deneo-Pelegrini (1997) [52]	175	Diet	↓	0.4 (0.2–0.8)
Kristal (1999) [57]	697	Suppl.	NS	0.77 (0.57–1.04)
<i>Cohort studies</i>				
Daviglus (1996) [50]	132	Diet	NS	1.03 (0.59–1.6)
Eichholzer (1999) [54]	30	Plasma	NS	0.93 (0.31–2.78) ²
NS = Statistically not significant; ↓ = statistically significant inverse association.				
¹ Comparing extreme categories, low = reference; ² high = reference.				

case-control study in Japan which showed that consumption of vegetables containing >600 µg carotin/100 g was associated with a decreased prostate cancer risk [39, 40]. An inverse correlation between higher carotene intake and prostate cancer risk was also observed in a case-control study in England [19]. These findings contrast to the results obtained in a recent case-control study by Norrish et al. [24] and in the Lutheran Brotherhood Cohort Study [41].

Most notable, the effects of β-carotene on prostate cancer risk have also been addressed in large interventional studies. The CARET trial (30 mg β-carotene/day) and the Physicians' Health Study (PHS) I (50 mg β-carotene every second day) could not demonstrate a beneficial effect of β-carotene on prostate cancer incidence [42–44]. However, in the PHS I trial those individuals with initial low plasma levels of β-carotene had a statistically significant 32% risk reduction after subsequent β-carotene supplementation as compared to those individuals not receiving β-carotene [43]. This contrast to a nonsignificant 33% increase in prostate cancer incidence in the individuals with high pre-study β-carotene plasma levels.

These findings could explain the observations made in the ATBC interventional trial including 29,133 male smokers aged 50–69 years, where the treatment group received 20 mg β-carotene/day alone or in combination with α-tocopherol (50 mg) [45–48]. In this group an increased incidence (23%) and mortality (15%) from prostate cancer was observed. Since no pre-study examination of the plasma levels was performed, a possible interaction of low or high β-carotene levels could not be disclosed.

In summary, the effects of β-carotene or total vitamin A on the development of prostate cancer remain inconclusive. There is even some evidence that uncon-

trolled β-carotene intake could be hazardous. However, so far, it appears that supplementation of β-carotene could be beneficial in a distinct group of men with low β-carotene plasma levels.

Vitamin C

The influence of vitamin C on prostate cancer has been studied in a number of case-control studies and cohort trials [49–57] (table 4). In the majority of these studies no evidence of an inverse association between vitamin C intake or vitamin C plasma concentrations on the incidence or mortality of prostate cancer was noted. A borderline significant inverse association was observed with daily supplemented use only in the case-control study by Kristal et al. [57]. It was suggested that higher than dietary amounts of vitamin C might be necessary for a protective effect. Although the results of prospective interventional trials are not yet available, it appears unlikely that vitamin C may play a major role in prostate cancer prevention.

Vitamin D

Vitamin D plays a crucial role in osseous metabolism and vitamin D deficiency or vitamin D receptor disorders are correlated with osteomalacia. Furthermore, it has been speculated that decreased vitamin D serum levels could be responsible for the increased risk in Black US citizens developing prostate cancer. Peehl et al. [35, 36] studied the antiproliferative effects of vitamin D₃ in primary cultures of prostatic tissues derived from prostate cancer patients. The antiproliferative effects of vitamin D₃ were more pronounced in prostate cancer cells as compared to prostate fibroblasts. A decreased expression of vitamin D receptors in fibroblasts was suggested to be responsible for this observation. Further investigation of the mechanism

Table 5. Summary of trials investigating the influence of serum vitamin D and calcium intake on the incidence of prostate cancer

Reference	Cases	Measure	Ass.	RR/OR (95% CI) ¹
<i>Case-control</i>				
Vlajinac (1997) [56]	101	Dietary calcium	↓	0.37 (0.14–0.99)
Chan (1998) [58, 59]	526	Dietary calcium	↑	1.9 (1.2–3.0)
Kristal (1999) [57]	697	Suppl. dietary calcium	NS	1.04 (0.61–1.78)
Ahonen (2000) [60]	149	Serum (vitamin D)	↓	1.7
<i>Cohort studies or nested case-control studies</i>				
Nomura (1988) [61]	136	Serum	NS	1.0 (0.5–2.1)
Giovannucci (1998) [63]	423	Diet (calcium intake)	↑	2.97 (1.6–5.5)
Schuurman (1999) [64]	642	Diet (calcium intake)	NS	n.r.
NS = Statistically not significant; ↓ = Statistically significant inverse association; ↑ = statistically significant direct association.				
¹ Comparing extreme categories, low = reference; n.r. = not reported.				

underlying the antiproliferative effects showed that growth inhibition was mediated both by androgen-dependent and androgen-independent pathways [37].

The relation between the intake of dairy products and prostate cancer risk has been investigated in a population-based case-control study by Chan et al. [58]. The investigation of 526 cases and 536 controls showed calcium intake to be an independent predictor of prostate cancer, and specifically for metastatic tumors. Since calcium consumption is supposed to lower vitamin D serum levels, the correlation between vitamin D metabolite serum levels and prostate cancer was studied in a nested case-control study in a Japanese-American population including 136 cases and 136 matched controls [61]. No difference between vitamin D serum levels in the two groups was observed. However, individuals with low vitamin D serum levels were under-represented which may hide an existing difference (table 5).

The investigation of stored sera collected from over 250,000 individuals through the years 1964–1971 showed significantly lower vitamin D serum levels in older patients (>57 years) with prostate cancer as compared to controls without prostate cancer [62]. Similarly, Ahonen et al. [60] reported a correlation between low vitamin D serum levels and subsequent development of prostate cancer in a Finnish study including 19,000 healthy men. However, in this trial low vitamin D serum levels were specifically linked to aggressive tumors in younger patients.

Other studies aiming at the relation between vitamin D receptor polymorphism and prostate cancer risk yielded controversial results. While a heterozygosity or homozygosity for the absence of the Bsm I restriction

site was associated with significant decrease of the prostate cancer risk in a Japanese population [65], this finding was not confirmed in a case-control study nested in the Physicians' Health Study [66].

So far, only phase I/II trials on a therapeutical use of vitamin D have been published. Gross et al. [67] demonstrated a significant decrease of serum PSA in 6 of 7 patients with early recurrent prostate cancer receiving 0.5–2.5 µg vitamin D₃/day for 6–15 months. Van Veldhuizen et al. [68] investigated vitamin D replacement therapy in patients with hormone-refractory metastatic prostate cancer. Four of the 16 patients had improvements in pain scores and 6 patients had improved muscle strength measurements. Notably, 7 of the 16 patients had decreased baseline vitamin D levels.

In summary, the role of vitamin D deficiency in the development of prostate cancer remains controversial. Despite the influence of high-dose vitamin D treatment on serum PSA, more detailed analyses are required to accept this approach as a viable therapeutic option.

Vitamine E (α-Tocopherol)

Vitamin E is a potent intracellular antioxidant with demonstrable antitumoral properties in several cancer models including sarcomas and gastrointestinal tumors [69–71]. It has been hypothesized that the antioxidant properties of vitamin E may reduce the dietary fat-induced oxidative stress and thus subsequently affect tumor cell growth.

Based on these investigations and further epidemiologic evidence that vitamin E could also have a preventive effect in prostate cancer, this question has been addressed in more detail in a variety of epidemiologi-

Table 6. Summary of trials investigating the influence of vitamin E on the incidence or mortality of prostate cancer

Reference	Cases	Measure	Ass.	RR/OR (95% CI) ¹
<i>Case-control</i>				
Rohan (1995) [53]	207	Diet	NS	n.r.
Andersson (1996) [72]	526	Diet	NS	0.91 (0.64–1.28)
Vlajinac (1997) [56]	101	Diet	↓	0.15 (0.05–0.53)
Deneo-Pelegrini (1999) [52]	175	Diet	↓ ptrend	0.6 (0.3–1.1)
Hayes (1999) [23]	932	Diet	NS	n.r. ²
Tzonou (1999) [25]	320	Diet	↓	0.53 (0.30–0.94)
Kristal (1999) [57]	697	Suppl.	NS	0.76 (0.54–1.08)
<i>Cohort studies or nested case-control studies</i>				
Knekt (1988) [73]	26	Serum	NS	0.42 (0.10–1.71)
Hsing (1990) [41]	103	Serum	NS	1.00 (0.37–2.68)
Nomura (1997) [29]	142	Serum	NS	0.9 (0.7–2.9)
Chan (1999/2000) [58, 59]	1,896	Suppl.	NS	1.07 (0.95–1.20)
Gann (1999) [31]	578	Plasma	NS	0.64 (0.38–1.07)
Eichholzer (1999) [54]	30	Plasma	NS	0.76 (0.25–2.37) ²
NS = Statistically not significant; ↓ = statistically significant inverse association.				
¹ Comparing extreme categories, low = reference; ² high = reference; n.r. = not reported.				

Table 7. Summary of trials investigating the influence of vitamin E on the incidence and mortality of prostate cancer in smokers

Reference	Cases	Measure	Ass.	RR/OR (95% CI) ¹
<i>Cohort studies or nested case-control studies</i>				
Knekt (1988) [73]	115 ²	Serum	NS	0.74 (0.44–1.26)
Chan (1999) [74]	55	Suppl.	↓	0.47 (0.24–0.92)
Gann (1999) [31]	578	Plasma	↓	0.51 (0.26–0.98)
Eichholzer (1999) [54]	30	Plasma	↓	3.26 (1.27–8.35) ²
<i>Interventional studies</i>				
ATBC [45–48]	62	50 mg	↓	0.59 (0.35–0.99)
NS = Statistically not significant; ↓ = statistically significant inverse association.				
¹ Comparing extreme categories, low = reference; ² High = reference.				

cal studies (table 6). However, although several case-control studies provided some evidence for a decreased incidence of prostate cancer in populations consuming a vitamin E-rich diet, this finding was not confirmed in subsequent cohort studies [5, 45–48].

Stratification for several risk factors in the Basel Cohort Study, however, showed that smokers with low vitamin E plasma levels had a significantly increased risk of prostate cancer mortality [54]. Reanalysis of the respective data from several other studies were mostly in concordance with this finding (table 7). In the ATBC interventional trial this hypothesis could be confirmed [45, 47]. Subjects receiving α -tocopherol (50 mg daily) for 5–8 years had a 32% decrease in the incidence and a 41% reduction in mortality from prostate cancer.

In summary, results of epidemiological studies do not support a general protective effect of vitamin E

against prostate cancer. However, results of four distinct populations, the US physicians' study, the Finnish ATBC trial, the Swiss Basel Study and the US health professionals' blood investigations, suggest that vitamin E may provide benefits against advanced prostate cancer or prostate cancer mortality among smokers or recent ex-smokers, but not among nonsmokers [31, 45–48, 54, 74].

Selenium

The inhibitory effect of selenomethionine, the predominant form of selenium in dietary supplements on the growth of prostate cancer cells and other tumor cell lines in vitro, has recently been studied by Redman et al. [75]. They found tumor cells to be approximately

Table 8. Summary of trials investigating the influence of dietary fat on the incidence of prostate cancer

Reference	Cases	Ass.	RR/OR (95% CI)
<i>Case-control studies</i>			
West (1991) [51]	679	↑	2.9 (1.08–8.4)
Ghadirian (1996) [83]	232	NS	n.r.
Vlajinac (1997) [56]	101	↑	3.6 (1.0–12.8)
Meyer (1997) [84–86]	215	↑	2.67 (??)
Lee (1998) [87]	133	↑	3.6 (1.8–7.2)
Deneo-Pelegrini (1999) [52]	175	↑Trend	1.8 (0.9–3.4)
Tzonou (1999) [25]	320	↑	n.r.
Ramon (2000) [55]	217	↑	2.0 (1.2–3.2)
<i>Cohort studies or nested case-control studies</i>			
Severson (1989) [17]	174	NS	0.91 (0.64–1.28)
Veierod (1997) [88]	72	NS	1.3 (0.6–2.8)
LeMarchand (1994) [89]	198	↑	1.6 (1.0–2.4)
Hayes (1999) [23]	932		
Total PCA		NS/↑ ²	1.1–2.1
Advanced PCA		↑/↑ ²	2.2–4.2
Schuurman (1999) [90]	642	NS	1.1

NS = Statistically not significant; ↑ = statistically significant direct association.
¹ Comparing extreme categories, low = reference; n.r. = not reported.
² White/Black US citizens.

1,000 times more sensitive as compared to normal diploid fibroblasts.

These observations match with data from a longitudinal case-control study on 111 subjects developing cancer within 5 years after obtaining blood samples for determination of serum selenium levels and compared the results with the levels of 210 control individuals not developing cancer within the same period [76]. The mean selenium levels of the cancer patients was significantly lower as compared with the controls. A similar study design was chosen in a prospective case-control study by Criqui et al. [77]. Baseline levels of plasma selenium, retinol and retinol-binding protein were compared in 136 case patients who subsequently died from cancer and 238 matched controls. Serum selenium was lower in case patients with gastrointestinal or prostate cancer, however, without reaching significance.

Hardell et al. [78] reported on a case-control study comparing selenium plasma levels from 164 patients with prostate cancer with 152 patients with benign prostatic hyperplasia. In those patients not taking supplementary selenium, the selenium levels were significantly lower in the cancer patients as compared to the controls. Yoshizawa et al. [79] conducted a nested case-control study within a large cohort study including

51,529 health professionals. Selenium levels were determined in toenail clippings. After controlling for several potentially interfering factors, higher selenium intake was found to be associated with a reduced risk for advanced prostate cancer.

For those patients enrolled in the ATBC study, baseline levels for several parameters were obtained [48]. No significant differences were reported for the incidence of prostate cancer and serum selenium levels. In an another interventional trial, Combs et al. [80] randomized 1,312 patients with a history of basal/squamous cell carcinoma in a double-blind fashion to either receive 200 µg selenium/day or placebo. While recurrence of skin cancer was not significantly reduced, total mortality and cancer mortality from all cancers, and in particular incidence and mortality from prostate cancer (secondary endpoints of the trial), were significantly reduced [81].

Dietary Fat

A high-fat diet has frequently been associated with an increased risk of prostate cancer. This aspect has been studied in in-vivo experiments where the growth of LNCaP cells in nude mice was investigated. Tumor

growth and serum PSA were significantly decreased in those animals receiving a low-fat diet as compared to control mice receiving an isocaloric high-fat diet [5].

Several aspects of increased fat intake, i.e. high serum cholesterol, total fat intake, the intake of saturated or unsaturated fatty acids or animal fat consumption have been studied in a variety of case-control or cohort studies (table 8). In a recent review addressing this topic, 11 out of 14 case-control trials yielded a significant correlation between fat consumption and prostate cancer risk [5]. Furthermore, 4 of the 5 cohort studies cited including nearly 100,000 individuals reported a relative risk ranging from 1.8 to 2.4 for people with high-fat intake. Even a correlation between fat consumption and high-grade/high-stage prostate cancer and prostate cancer mortality has been suggested by several studies [5, 82]. Meyer et al. [84–86] showed that a high intake of saturated fatty acids in patients with prostate cancer was significantly associated with tumor progression and mortality. In contrast to most earlier studies, these data were adjusted for total calorie intake.

These observations are further supported by studies reporting a correlation between overweight (body mass index) and the prostate cancer risk [5, 30, 91, 92]. In contrast to the obvious-tumor-promoting effects of saturated fats especially from animal sources, the intake of fat derived from plants or ω -3 fatty acid appears not relate to an increased risk developing prostate cancer [5, 93, 94].

Conclusions

Large autopsy studies suggest that the incidence latent prostate cancer is similar worldwide. This observation contrasts to the regionally different incidence of overt or clinically relevant tumors. Therefore, already a long time ago an environmental influence on the progression of microscopic to clinically relevant prostate cancer was postulated. This hypothesis was supported by migration studies demonstrating the increase of prostate cancer incidence in populations moving from areas with a low incidence in the Far East to areas with high incidence in the US [1, 2].

Throughout the last two decades a significant number of studies on the impact of nutritional compounds on the development but also on the course of disease of prostate cancer have been published. Considering the general question regarding an effect of nutrition on prostate cancer there is – despite several contradictory

Table 9. Factors influencing the result of trials investigating the impact of nutrition on prostate cancer

Validity of studies (case-control, cohort studies, interventional trials)
Problem of follow-up interval in the interventional and most cohort studies
Reliability of questionnaires
Hereditary factors
Data from different ethnical populations
Correlation between nutrition and lifestyle
Interaction between several nutritional compounds

reports – increasing evidence supporting this hypothesis [95]. However, when it comes to more detailed questions aiming at the nature of relevant nutritional compounds, instead of clear conclusions more and more open questions remain.

These contradictory observations are better understood in the light of the complexity of the problem. So far, multiple factors potentially influencing the results of the respective trials have been identified (table 9).

Hypotheses of actual trials are frequently based upon results obtained in studies of questionable quality. Most results discussed in numerous review articles on this topic have been obtained by case-control studies. However, the validity of this type of trial is limited since selection bias is immanent and may be responsible for conflicting results [96]. In contrast, the impact of cohort studies is certainly superior, but high costs and long follow-up intervals required in general represent significant obstacles to launch this type of study. The most valuable information can be obtained through interventional trials, however until today only very few trials of this type have been generated to study the role of nutrition on prostate cancer [42–48, 97].

The validity of questionnaires is another problem that may distort the results of those trials relaying on self-reports of individuals. An investigation on the reliability of self-reports demonstrated a high level of inaccuracy specifically if non-life-threatening issues were assessed [98–100]. This has led to the inclusion of objective parameters through the analysis of serum, tissue, saliva and toenails in more recent trials [81, 101, 102].

Furthermore, the fact that hereditary factors also affect the results of epidemiologic studies on prostate cancer [103] and, furthermore, that data from different

ethnic populations yield different results further impedes the interpretation of the reports [104]. However, even within ethnically homogeneous populations, the correlation between nutrition and life-style may be confounded by genetic factors. Cross-national (meta)-analyses could be one option to address this problem but the quality of the different trials included in this type of study is hard to control for [105].

Apart from these factors that could be theroretically taken into account designing a trial on nutrition in prostate cancer, the interaction between the different nutritional compounds remains a key problem in population-based trials. Considering the putative beneficial effects of tomatoes and tomato products, it is nearly impossible to identify which of the multiple compounds or which combination of compounds may be responsible for beneficial effects. And even the administration of a single compound – e.g. a vitamin – may be influenced by other interacting nutritional or lifestyle factors.

However, these problems should not discourage the scientific community to conduct further well-designed trials on this import issue but should caution inadequate expectations and overinterpretation of the results of single trials. Today the concept of an impact of nutrition on prostate cancer appears well established but the details still require further characterization. Primary and secondary as well as tertiary prevention need to be addressed [97]. Considering the noninvasiveness of this approach and the putative benefits, a thorough investigation of this issue appears worthwhile and certainly justifies the efforts invested in this topic worldwide [97, 106, 107].

Acknowledgements

This manuscript is dedicated to Prof. R. Ackermann on the occasion of his 60th birthday.

The secretarial assistance of Mrs. S. Zaun is gratefully acknowledged.

References

- Muir CS, Nectoux J, Staszewski J: The epidemiology of prostatic cancer: Geographical distribution and timetrends. *Acta Oncol* 1991;30: 133–140.
- Haenszel W, Kurihara M: Mortality from cancer and other diseases among Japanese in the United States. *J Natl Cancer Inst* 1968;40:43–68.
- Kelloff GJ, Lieberman R, Steele VE, Boone CW, Lubet RA, Kopelovitch L, Malone WA, Crowell JA, Sigman CC: Chemoprevention of prostate cancer: Concepts and strategies. *Eur Urol* 1999;35:342–345.
- Kelloff GJ, Crowell JA, Steele VE, Lubet RA, Boone CW, Malone WA, Hawk ET, Lieberman R, Lawrence JA, Kopelovich L, Ali I, Viner JL, Sigman CC: Progress in cancer chemoprevention. *Ann NY Acad Sci* 1999; 889:1–13.
- Fair WR, Fleshner NE, Heston W: Fleshner NE, Heston W: Cancer of the prostate: A nutritional disease? *Urology* 1997;50:840–848.
- Gann PH, Hennekens CH, Ma J, Longcope C, Stampfer MJ: Prospective study of sex hormone levels and risk of prostate cancer. *J Natl Cancer Inst* 1996;88:1118–1126.
- Nomura AM, Stemmermann GN, Chyou PH, Henderson BE, Stanczyk FZ: Serum androgens and prostate cancer. *Cancer Epidemiol Biomarkers Prev* 1996;5:621–625.
- Dorgan JF, Albanes D, Virtamo J, Heinonen OP, Chandler DW, Galmarini M, McShane LM, Barrett MJ, Tangrea J, Taylor PR: Relationships of serum androgens and estrogens to prostate cancer risk: Results from a prospective study in Finland. *Cancer Epidemiol Biomarkers Prev* 1998;7:1069–1074.
- Graham S, Haughey B, Marshall J, Priore R, Byers T, Rzepka T, Mettlin C, Pontes JE: Diet in the epidemiology of carcinoma of the prostate gland. *J Natl Cancer Inst* 1983;70: 687–692.
- Aronson WJ, Tymchuk CN, Elashoff RM, McBride WH, McLean C, Wang H, Heber D: Decreased growth of human prostate LNCaP tumors in SCID mice fed a low-fat, soy protein diet with isoflavones. *Nutr Cancer* 1999; 35:130–136.
- Griffiths K, Denis L, Turkes A, Morton MS: Possible relationship between dietary factors and pathogenesis of prostate cancer. *Int J Urol* 1998;5:195–213.
- Morton MS, Matos-Ferreira A, Abranches-Monteiro L, Correia R, Blacklock N, Chan PS, Cheng C, Lloyd S, Chieh-Ping W, Griffiths K: Measurement and metabolism of isoflavonoids and lignans in the human male. *Cancer Lett* 1997;114:145–151.
- Adlercreutz H, Markkanen H, Watanabe S: Plasma concentrations of phyto-oestrogens in Japanese men. *Lancet* 1993;342:1209–1210.
- Strom SS, Yamamura Y, Duphorne CM, Spitz MR, Babaian RJ, Pillow PC, Hursting SD: Phytoestrogen intake and prostate cancer: A case-control study using a new database. *Nutr Cancer* 1999;33:20–25.
- Kolonel LN, Hankin JH, Whittemore AS, Wu AH, Gallagher RP, Wilkens LR, John EM, Howe GR, Dreon DM, West DW, Paffenbarger RS Jr: Vegetables, fruits, legumes and prostate cancer: A multiethnic case-control study. *Cancer Epidemiol Biomarkers Prev* 2000;9:795–804.
- Jain MG, Hislop GT, Howe GR, Ghadirian P: Plant foods, antioxidants, and prostate cancer risk: Findings from case-control studies in Canada. *Nutr Cancaer* 1999;34:173–184.
- Severson RK, Nomura AM, Grove JS, Stemmermann GN: A prospective study of demographics, diet, and prostate cancer among men of Japanese ancestry in Hawaii. *Cancer Res* 1989;49:1857–1860.
- Jacobsen BK, Knutsen SF, Fraser GE: Does high soy milk intake reduce prostate cancer incidence? The Adventist Health Study. *Cancer Causes Control* 1998;9:553–557.
- Key TJ, Fraser GE, Thorogood M, Appleby PN, Beral V, Reeves G, Burr ML, Chang-Claude J, Frentzel-Beyme R, Kuzma JW, Mann J, McPherson K: Mortality in vegetarians and nonvegetarians: Detailed findings from a collaborative analysis of five prospective studies. *Am J Clin Nutr* 1999;70(suppl):516S–524S.
- Agarwal S, Rao AV: Tomato lycopene and its role in human health and chronic diseases. *Can Med Assoc J* 2000;163:739–744.
- Le Marchand L, Hankin JH, Kolonel LN, Wilkens LR: Vegetable and fruit consumption in relation to prostate cancer risk in Hawaii: A reevaluation of the effect of dietary beta-carotene. *Am J Epidemiol* 1991;133:215–219.
- Key TJ, Silcocks PB, Dabey GK, Appleby PN, Bishop DT: A case-control study of diet and prostate cancer. *Br J Cancer* 1997;76:678–687.

- 23 Hayes RB, Ziegler RG, Gridley G, Swanson C, Greenberg RS, Swanson GM, Schoenberg JB, Silverman DT, Brown LM, Pottern LM, Liff J, Schwartz AG, Fraumeni JF Jr, Hoover RN: Dietary factors and risks for prostate cancer among Blacks and Whites in the United States. *Cancer Epidemiol Biomarkers Prev* 1999; 8:25–34.
- 24 Norrish AE, Jackson RT, Sharpe SJ, Skeaff CM: Prostate cancer and dietary carotenoids. *Am J Epidemiol* 2000;151:119–123.
- 25 Tzonou A, Signorello LB, Lagiou P, Wu J, Trichopoulos D, Trichopoulos A: Diet and cancer of the prostate: A case-control study in Greece. *Int J Cancer* 1999;80:704–708.
- 26 Mills PK, Beeson WL, Phillips RL, Fraser GE: Cohort study of diet, lifestyle, and prostate cancer in Adventist men. *Cancer* 1989;64:598–604.
- 27 Hsing AW, Comstock GW, Abbey H, Polk BF: Serologic precursors of cancer. Retinol, carotenoids, and tocopherol and risk of prostate cancer. *J Natl Cancer Inst* 1990;82: 941–946.
- 28 Giovannucci E, Ascherio A, Rimm EB, Stampfer MJ, Colditz GA, Willett WC: Intake of carotenoids and retinol in relation to risk of prostate cancer. *J Natl Cancer Inst* 1995; 87:1767–1776.
- 29 Nomura AM, Stemmermann GN, Lee J, Craft NE: Serum micronutrients and prostate cancer in Japanese Americans in Hawaii. *Cancer Epidemiol Biomarkers Prev* 1997;6:487–491.
- 30 Cerhan JR, Torner JC, Lynch CF, Rubenstein LM, Lemke JH, Cohen MB, Lubaroff DM, Wallace RB: Association of smoking, body mass, and physical activity with risk of prostate cancer in the Iowa 65+ Rural Health Study (United States). *Cancer Causes Control* 1997; 8:229–238.
- 31 Gann PH, Ma J, Giovannucci E, Willett W, Sacks FM, Hennekens CH, Stampfer MJ: Lower prostate cancer risk in men with elevated plasma lycopene levels: Results of a prospective analysis. *Cancer Res* 1999;59:1225–1230.
- 32 Giovannucci E: Tomatoes, tomato-based products, lycopene, and cancer: Review of the epidemiologic literature. *J Natl Cancer Inst* 1999;91:317–331.
- 33 Fleshner N, Fair W, Huryk R, Heston WDW: Vitamin E inhibits the high-fat diet-promoted growth of established human LnCaP tumors in nude mice. *J Urol* 1999;161:1651–1654.
- 34 Schwartz GG, Hill CC, Oeler TA, Becich MJ, Bahnson RR: 1,25-Dihydroxy-16-ene-23-yne-vitamin D3 and prostate cancer cell proliferation in vivo. *Urology* 1995;46:365–369.
- 35 Peehl DM, Skowronski RJ, Leung GK, Wong ST, Stamey TA, Feldman D: Antiproliferative effects of 1,25-dihydroxyvitamin D3 on primary cultures of human prostatic cells. *Cancer Res* 1994;54:805–810.
- 36 Peehl DM, Wong ST, Stamey TA: Vitamin A regulates proliferation and differentiation of human prostatic epithelial cells. *Prostate* 1993; 23:69–78.
- 37 Zaho XY, Peehl DM, Navone NM, Feldman D: 1 α ,25-Dihydroxyvitamin D3 inhibits prostate cancer cell growth by androgen-dependent and androgen-independent mechanisms. *Endocrinology* 2000;141:2548–2556.
- 38 Williams AW, Boileau TW, Zhou JR, Clinton SK, Erdman JW Jr: Beta-carotene modulates human prostate cancer cell growth and may undergo intracellular metabolism to retinol. *J Nutr* 2000;130:728–732.
- 39 Ohno Y, Yoshida O, Oishi K, Okada K, Yamabe H, Schroeder FH: Dietary beta-carotene and cancer of the prostate: A case-control study in Kyoto, Japan. *Cancer Res* 1988;48: 1331–1336.
- 40 Oishi K, Okada K, Yoshida O, Yamabe H, Ohno Y, Hayes RB, Schroeder FH: A case-control study of prostatic cancer with reference to dietary habits. *Prostate* 1988;12:179–190.
- 41 Hsing AW, McLaughlin JK, Schuman LM, Bjelke E, Gridley G, Wacholder S, Chien HT, Blot WJ: Diet, tobacco use, and fatal prostate cancer: Results from the Lutheran Brotherhood Cohort Study. *Cancer Res* 1990;50: 6836–6840.
- 42 Omenn GS, Goodman GE, Thornquist MD, Balmes J, Cullen MR, Glass A, Keogh JP, Meyskens FL Jr, Valanis B, Williams JH Jr, Barnhart S, Cherniack MG, Brodtkin CA, Hammer S: Risk factors for lung cancer and for intervention effects in CARET, the Beta-Carotene and Retinol Efficacy Trial. *J Natl Cancer Inst* 1996;88:1550–1559.
- 43 Cook NR, Le IM, Manson JE, Buring JE, Hennekens CH: Effects of beta-carotene supplementation on cancer incidence by baseline characteristics in the Physicians' Health Study. *Cancer Causes Control* 2000;11:617–626.
- 44 Cook NR, Stampfer MJ, Ma J, Manson JE, Sacks FM, Buring JE, Hennekens CH: Beta-carotene supplementation for patients with low baseline levels and decreased risks of total and prostate carcinoma. *Cancer* 1999;86:1783–1792.
- 45 Albanes D, Heinonen OP, Huttunen JK, Taylor PR, Virtamo J, Edwards BK, Haapakoski J, Rautalahti M, Hartman AM, Palmgren J, et al: Effects of alpha-tocopherol and beta-carotene supplements on cancer incidence in the Alpha-Tocopherol Beta-Carotene Cancer Prevention Study. *Am J Clin Nutr* 1995;62(suppl):1427–1430.
- 46 Alpha-Tocopherol, Beta-Carotene Cancer Prevention Study Group: The effect of vitamin E and beta-carotene on the incidence of lung cancers and other cancers in male smokers. *New Engl J Med* 1994;330:1029.
- 47 Heinonen OP, Albanes D, Virtamo J, Taylor PR, Huttunen JK, Hartman AM, Haapakoski J, Malila N, Rautalahti M, Ripatti S, Maenpaa H, Teerenhovi L, Koss L, Virolainen M, Edwards BK: Prostate cancer and supplementation with alpha-tocopherol and beta-carotene: Incidence and mortality in a controlled trial. *J Natl Cancer Inst* 1998;90: 440–446.
- 48 Hartman TJ, Albanes D, Pietinen P, Hartman AM, Rautalahti M, Tangrea JA, Taylor PR: The association between baseline vitamin E, selenium, and prostate cancer in the alpha-tocopherol, beta-carotene cancer prevention study. *Cancer Epidemiol Biomarkers Prev* 1998;7:335–340.
- 49 Kolonel LN, Hinds MW, Nomura AM, Hankin JH, Lee J: Relationship of dietary vitamin A and ascorbic acid intake to the risk for cancers of the lung, bladder, and prostate in Hawaii. *Natl Cancer Inst Monogr* 1985;69:137–142.
- 50 Daviglus ML, Dyer AR, Persky V, Chavez N, Drum M, Goldberg J, Liu K, Morris DK, Shekelle RB, Stamler J: Dietary beta-carotene, vitamin C, and risk of prostate cancer: Results from the Western Electric Study. *Epidemiology* 1996;7:472–477.
- 51 West DW, Slattery ML, Robinson LM, French TK, Mahoney AW: Adult dietary intake and prostate cancer risk in Utah: A case-control study with special emphasis on aggressive tumors. *Cancer Causes Control* 1991;2:85–94.
- 52 Deneo-Pellegrini H, De Stefani E, Ronco A, Mendilaharsu M: Foods, nutrients and prostate cancer: A case-control study in Uruguay. *Br J Cancer* 1999;80:591–597.
- 53 Rohan TE, Howe GR, Burch JD, Jain M: Dietary factors and risk of prostate cancer: A case-control study in Ontario, Canada. *Cancer Causes Control* 1995;6:145–154.
- 54 Eichholzer M, Stahelin HB, Ludin E, Bernasconi F: Smoking, plasma vitamins C, E, retinol, and carotene, and fatal prostate cancer: Seventeen-year follow-up of the prospective Basel study. *Prostate* 1999;38:189–198.
- 55 Ramon JM, Bou R, Romea S, Alkiza ME, Jacas M, Ribes J, Oromi J: Dietary fat intake and prostate cancer risk: A case-control study in Spain. *Cancer Causes Control* 2000;11:679–685.
- 56 Vlainjac HD, Marinkovic JM, Ilic MD, Kocov NI: Diet and prostate cancer: A case-control study. *Eur J Cancer* 1997;33: 101–107.
- 57 Kristal AR, Stanford JL, Cohen JH, Wicklund K, Patterson RE: Vitamin and mineral supplement use is associated with reduced risk of prostate cancer. *Cancer Epidemiol Biomarkers Prev* 1999;8:887–892.
- 58 Chan JM, Giovannucci E, Andersson SO, Yuen J, Adami HO, Wolk A: Dairy products, calcium, phosphorus, vitamin D, and risk of prostate cancer. *Cancer Causes Control* 1998; 9:559–566.
- 59 Chan JM, Pientinen P, Virtanen M, Malila N, Tangrea J, Albanes D, Virtamo J: Diet and prostate cancer risk in a cohort of smokers, with a specific focus on calcium and phosphorus. *Cancer Causes Control* 2000;11:859–867.
- 60 Ahonen MH, Tenkanen L, Teppo L, Hakama M, Tuohimaa P: Prostate cancer risk and prediagnostic serum 25-hydroxyvitamin D levels. *Cancer Causes Control* 2000;11:847–852.
- 61 Nomura AM, Stemmermann GN, Lee J, Kolonel LN, Chen TC, Turner A, Holick MF: Serum vitamin D metabolite levels and the subsequent development of prostate cancer. *Cancer Causes Control* 1998;9:425–432.
- 62 Corder EH, Guess HA, Hulka BS, Friedman GD, Sadler M, Vollmer RT, Lobaugh B, Drezner MK, Vogelstein JH, Orentreich N: Vitamin D and prostate cancer: A prediagnostic study with stored sera. *Cancer Epidemiol Biomarkers Prev* 1993;2:467–472.
- 63 Giovannucci E, Rimm EB, Wolk A, Ascherio A, Stampfer MJ, Colditz GA, Willett WC: Calcium and fructose intake in relation to risk of prostate cancer. *Cancer Res* 1998;58:442–447.
- 64 Schuurman AG, van den Brandt PA, Dorant E, Goldbohm RA: Animal products, calcium and protein and prostate cancer risk in The Netherlands Cohort Study. *Br J Cancer* 1999;80: 1107–1113.

- 65 Habuchi T, Suzuki T, Sasaki R, Wang L, Sato K, Satoh S, Akao T, Tsuchiya N, Shimoda N, Wada Y, Koizumi A, Chihara J, Ogawa O, Kato T: Association of vitamin D receptor gene polymorphism with prostate cancer and benign prostatic hyperplasia in a Japanese population. *Cancer Res* 2000;60:305–308.
- 66 Ma J, Stampfer MJ, Gann PH, Hough HL, Giovannucci E, Kelsey KT, Hennekens CH, Hunter DJ: Vitamin D receptor polymorphisms, circulating vitamin D metabolites, and risk of prostate cancer in United States physicians. *Cancer Epidemiol Biomarkers Prev* 1998;7:385–399.
- 67 Gross C, Stamey T, Hancock S, Feldman D: Treatment of early recurrent prostate cancer with 1,25-dihydroxyvitamin D₃. *J Urol* 1998;159:2035–2039.
- 68 Van Veldhuizen PJ, Taylor SA, Williamson S, Drees BM: Treatment of vitamin D deficiency in patients with metastatic prostate cancer may improve bone pain and muscle strength. *J Urol* 2000;163:187–190.
- 69 Das S: Vitamin E in the genesis and prevention of cancer. *Acta Oncol* 1994;33:615.
- 70 Vasavi H, Thangaraju M, Sachdanandam P: Effects of alpha-tocopherol on lipid peroxidation and antioxidant system in fibrosarcoma-bearing rats. *Mol Cell Biochem* 1994;131:125–129.
- 71 Shivapakur H: Inhibition of progression of aberrant crypt foci and colon tumor development by vitamin E and beta-carotene in rats on a high fat diet. *Cancer Lett* 1995;91:125.
- 72 Andersson SO, Wolk A, Bergstrom R, Giovannucci E, Lindgren C, Baron J, Adami HO: Energy, nutrient intake and prostate cancer risk: A population-based case-control study in Sweden. *Int J Cancer* 1996;68:716–722.
- 73 Knekt P, Aromaa A, Maatela J, Aaran RK, Nikkari T, Hakama M, Hakulinen T, Peto R, Saxen E, Teppo L: Serum vitamin E and risk of cancer among Finnish men during a 10-year follow-up. *Am J Epidemiol* 1988;127:28–41.
- 74 Chan JM, Stampfer MJ, Ma J, Rimm EB, Willett WC, Giovannucci EL: Supplemental vitamin E intake and prostate cancer risk in a large cohort of men in the United States. *Cancer Epidemiol Biomarkers Prev* 1999;8:893–899.
- 75 Redman C, Scott JA, Baines AT, Basye JL, Clark LC, Calley C, Rose D, Payne CM, Nelson MA: Inhibitory effect of selenomethionine on the growth of three selected human tumor cell lines. *Cancer Lett* 1998;125:103–110.
- 76 Willett WC, Polk BF, Morris JS, Stampfer MJ, Pressel S, Rosner B, Taylor JO, Schneider K, Hames CG: Prediagnostic serum selenium and risk of cancer. *Lancet* 1983;iii:130–134.
- 77 Criqui MH, Bangdiwala S, Goodman DS, Blaner WS, Morris JS, Kritchevsky S, Lippel K, Mebane I, Tyroler HA: Selenium, retinol, retinol-binding protein, and uric acid. Associations with cancer mortality in a population-based prospective case-control study. *Am Epidemiol* 1991;138:385–393.
- 78 Hardell L, Degerman A, Tomic R, Marklund SL, Bergfors M: Levels of selenium in plasma and glutathione peroxidase in erythrocytes in patients with prostate cancer or benign hyperplasia. *Eur J Cancer Prev* 1995;4:91–95.
- 79 Yoshizawa K, Willett WC, Morris SJ, Stampfer MJ, Spiegelman D, Rimm EB, Giovannucci E: Study of prediagnostic selenium level in toenails and the risk of advanced prostate cancer. *J Natl Cancer Inst* 1998;90:1219–1224.
- 80 Combs GF Jr, Clark LC, Turnbull BW: Reduction of cancer risk with an oral supplement of selenium. *Biomed Environ Sci* 1997;10:227–234.
- 81 Clark LC, Dalkin B, Krongrad A, Combs GF Jr, Turnbull BW, Slate EH, Witherington R, Herlong JH, Janosko E, Carpenter D, Borosso C, Falk S, Rounder J: Decreased incidence of prostate cancer with selenium supplementation: Results of a double-blind cancer prevention trial. *Br J Urol* 1998;81:730–734.
- 82 Rose DP, Boyer AP, Wynder EL: International comparison of mortality rates for cancer of the breast, ovary, prostate, and colon per capita food consumption. *Cancer* 1986;58:2363–2371.
- 83 Ghadirian P, Lacroix A, Maisonneuv P, Perret C, Drouin G, Perrault JP, Beland G, Rohan TE, Howe GR: Nutritional factors and prostate cancer: A case-control study of French Canadians in Montreal, Canada. *Cancer Causes Control* 1996;7:428–436.
- 84 Meyer F, Bairati I, Fradet Y, Moore L: Dietary energy and nutrients in relation to preclinical prostate cancer. *Nutr Cancer* 1997;29:120–126.
- 85 Meyer F, Bairati I, Shadmani R, Fradet Y, Moore L: Dietary fat and prostate cancer survival. *Cancer Causes Control* 1999;10:245–251.
- 86 Fradet Y, Meyer F, Bairati I, Shadmani R, Moore L: Dietary fat and prostate cancer progression and survival. *Eur Urol* 1999;35:388–391.
- 87 Lee MM, Wang RT, Hsing AW, Gu FL, Wang T, Spitz M: Case-control study of diet and prostate cancer in China. *Cancer Causes Control* 1998;9:545–552.
- 88 Veierod MB, Laake P, Thelle DS: Dietary fat intake and risk of prostate cancer: A prospective study of 25,708 Norwegian men. *Int J Cancer* 1997;73:634–638.
- 89 Le Marchand L, Kolonel LN, Wilkens LR, Myers BC, Hirohata T: Animal fat consumption and prostate cancer: A prospective study in Hawaii. *Epidemiology* 1994;5:276–282.
- 90 Schuurman AG, van den Brandt PA, Dorant E, Brants HA, Goldbohm RA: Association of energy and fat intake with prostate carcinoma risk: Results from The Netherlands Cohort Study. *Cancer* 1999;86:1019–1027.
- 91 Andersson SO, Wolk A, Bergstrom R, Adami HO, Engholm G, Englund A, Nyren O: Body size and prostate cancer: A 20-year follow-up study among 135,006 Swedish construction workers. *J Natl Cancer Inst* 1997;89:385–389.
- 92 Chyou PH, Nomura AM, Stemmermann GN: A prospective study of weight, body mass index and other anthropometric measurements in relation to site-specific cancers. *Int J Cancer* 1994;57:313–317.
- 93 Norrish AE, Jackson RT, Sharpe SJ, Skeaff CM: Men who consume vegetable oils rich in monounsaturated fat: Their dietary patterns and risk of prostate cancer. *Cancer Causes Control* 2000;11:609–615.
- 94 Norrish AE, Skeaff CM, Arribas GL, Sharpe SJ, Jackson RT: Prostate cancer risk and consumption of fish oils: A dietary biomarker-based case-control study. *Br J Cancer* 1999;81:1238–1242.
- 95 Blumenfeld AJ, Fleshner N, Casselman B, Trachtenberg J: Nutritional aspects of prostate cancer: A review. *Can J Urol* 2000;7:927–935.
- 96 Kim DJ, Gallagher RP, Hislop TG, Holowaty EJ, Howe GR, Jain M, McLaughlin JR, Teh CZ, Rohan TE: Premorbid diet in relation to survival from prostate cancer (Canada). *Cancer Causes Control* 2000;11:65–77.
- 97 Schröder FH, Krane R, Dijk MA, Blom JM, Tijburg LM, Weststrate JA, Dagnelie PC: Tertiary prevention of prostatic cancer by dietary intervention: Results of a randomized, double-blind, placebo-controlled cross-over study. *Eur Urol* 2000;37(suppl 2):24.
- 98 Bergmann MM, Byers T, Freedman DS, Mokdad A: Validity of self-reported diagnoses leading to hospitalization: A comparison of self-reports with hospital records in a prospective study of American adults. *Am J Epidemiol* 1998;147:969–977.
- 99 Neuhauser ML, Kristal AR, McLerran D, Patterson RE, Atkinson J: Validity of short food frequency questionnaires used in cancer chemoprevention trials: Results from the Prostate Cancer Prevention Trial. *Cancer Epidemiol Biomarkers Prev* 1999;8:721–725.
- 100 Freeman VL, Meydani M, Yong S, Pyle J, Wan Y, Arvizu-Durazo R, Liao Y: Prostatic levels of tocopherols, carotenoids, and retinol in relation to plasma levels and self-reported usual dietary intake. *Am J Epidemiol* 2000;151:109–118.
- 101 Clinton SK, Emenhiser C, Schwartz SJ, Bostwick DG, Williams AW, Moore BJ, Erdmann JW Jr: cis-trans lycopene isomers, carotenoids, and retinol in the human prostate. *Cancer Epidemiol Biomarkers Prev* 1996;5:823–833.
- 102 Rao AV, Fleshner N, Agarwal S: Serum and tissue lycopene and biomarkers of oxidation in prostate cancer patients: A case-control study. *Nutr Cancer* 1999;33:159–164.
- 103 Greenwald P: Cancer risk factors for selecting cohorts for large-scale chemoprevention trials. *J Cell Biochem Suppl* 1996;25:29–36.
- 104 Hanash KA, Al-Othaimen A, Kattan S, Lindstedt E, Al-Zahrani H, Merdad T, Peracha A, Kardar AH, Aslam M, Al-Akkad A: Prostatic carcinoma: A nutritional disease? Conflicting data from the Kingdom of Saudi Arabia. *J Urol* 2000;164:1570–1572.
- 105 Grant WB: An ecologic study of dietary links to prostate cancer. *Altern Med Rev* 1999;4:162–169.
- 106 Christen WG, Gaziano JM, Hennekens CH: Design of Physicians' Health Study II – A randomized trial of beta-carotene, vitamins E and C, and multivitamins, in prevention of cancer, cardiovascular disease, and eye disease, and review of results of completed trials. *Ann Epidemiol* 2000;10:125–134.
- 107 Thompson IM, Coltman CA Jr, Crowley J: Chemoprevention of prostate cancer: The Prostate Cancer Prevention Trial. *Prostate* 1997;33:217–221.